

COVID-19 VACCINATION SUMMARY TABLE

Product	Dose	6 Months – 4 Years	5 Years and Older
Pfizer-BioNTech (COMIRNATY) <i>Appropriate to mix products for boosters, but preference is for a Pfizer or Moderna product</i>	Primary Series	3 doses; 2 nd dose 3-8 weeks after 1 st dose and 3 rd dose at least 8 weeks after 2 nd dose <u>Up to Date</u> = 2 weeks after final dose of primary series Note: For people who are moderately or severely immunocompromised (have a weakened immune system), the first and second doses are separated by three weeks, and the second and third doses are separated by at least eight weeks.	2 doses 3-8 weeks apart <u>Fully Vaccinated</u> = 2 weeks after final dose of primary series Note: People who are moderately or severely immunocompromised (have a weakened immune system) should get a 3rd dose at least 4 weeks after their 2nd dose in the primary series.
	Booster 1	Not yet authorized for this age group	Given at least 5 months after final dose in primary series <u>Up to Date</u> = immediately after booster and not eligible for a second booster
	Booster 2	Not yet authorized for this age group	Given at least 4 months after first booster for <u>adults ages 50 years and older/those who are moderately or severely immunocompromised</u> <u>Up to Date</u> = immediately after booster if eligible

**Click on document to see Moderna and Johnson & Johnson information*

Product	Dose	6 Months – 17 Years	18 Years and Older
Moderna (Spikevax) <i>Appropriate to mix products for boosters, but preference is for a Pfizer or Moderna product</i>	Primary Series	2 doses 4-8 weeks apart <i>Up to Date = 2 weeks after final dose in primary series</i> Note: People who are moderately or severely immunocompromised (have a weakened immune system) should get a third dose – with the first and second doses separated by 4 weeks, and the second and third doses separated by at least 4 weeks.	2 doses 4-8 weeks apart <i>Fully Vaccinated = 2 weeks after final dose of primary series</i> Note: People who are moderately or severely immunocompromised (have a weakened immune system) should get a third dose – with the first and second doses separated by 4 weeks, and the second and third doses separated by at least 4 weeks.
	Booster 1	Not yet authorized for this age group	Given at least 5 months after final dose in primary series <i>Up to Date = immediately after booster</i>
	Booster 2	Not yet authorized for this age group	Given at least 4 months after first booster for <u>adults ages 50 years and older/those who are moderately or severely immunocompromised</u> <i>Up to Date = immediately after booster if eligible</i>

Product	Dose	18+ Years
Johnson & Johnson (Janssen) <i>Moderna or Pfizer preferred for primary series; boosters should be a Pfizer or Moderna product</i>	Primary Series	1 dose <u>Fully Vaccinated</u> = 2 weeks after primary dose Note: People who are moderately or severely immunocompromised (have a weakened immune system) should get a 2nd dose of Pfizer-BioNTech or Moderna at least 4 weeks after their 1st dose in the primary series.
	Booster 1 (Pfizer-BioNTech or Moderna product preferred)	Given at least 2 months after primary dose <u>Up to Date</u> = immediately after booster
	Booster 2 (Pfizer-BioNTech or Moderna product ONLY)	Given at least 4 months after first booster for <u>adults ages 50 years and older/those who are moderately or severely immunocompromised</u> <u>Up to Date</u> = immediately after booster if eligible Janssen vaccine can NOT be used for a second booster.

*Considerations:

People ages 5 through 64 years, and especially males ages 12 through 39 years, may consider getting the 2nd primary dose of an mRNA COVID-19 vaccine (Pfizer-BioNTech or Moderna; Pfizer-BioNTech only for children ages 5 through 17 years) 8 weeks after the 1st dose. A longer time between the 1st and 2nd primary doses may increase how much protection the vaccines offer, and further minimize the rare risk of heart problems, including myocarditis and pericarditis.

People ages 65 years and older, people more likely to get very sick from COVID-19, or anyone wanting protection due to high levels of community transmission should get the 2nd primary dose of Pfizer-BioNTech COVID-19 vaccine 3 weeks (or 21 days) after the 1st primary dose, or the 2nd primary dose of Moderna COVID-19 vaccine 4 weeks (or 28 days) after the 1st primary dose.