

COVID-9 VACCINATION SUMMARY TABLE

Pfizer-BioNTech

Age Group	Dose 1	Dose 2	Dose 3/Bivalent Booster
6 months – 4 years	Primary Series	Primary Series; given 3-8 weeks after 1 st dose	Primary Series; given at least 8 weeks after 2 nd dose <i>*the bivalent Pfizer-BioNTech product is used for this dose</i> <u>Up to Date</u> = 2 weeks after 3 rd dose
5 years – 17 years	Primary Series	Primary Series; given 3-8 weeks after 1 st dose	Updated Booster; given at least 2 months after 2 nd dose/last booster; can only be a Pfizer-BioNTech product if 5 years of age; can be either Pfizer-BioNTech or Moderna product if 6 years of age or older <u>Up to Date</u> = immediately after booster
18 years and older	Primary Series	Primary Series; given 3-8 weeks after 1 st dose	Updated Booster; given at least 2 months after 2 nd dose/last booster; can be either Pfizer-BioNTech or Moderna product <u>Up to Date</u> = immediately after booster

Moderna

Age Group	Dose 1	Dose 2	Bivalent Booster
6 months – 17 years	Primary Series	Primary Series; given 4-8 weeks after 1 st dose	Updated Booster; given at least 2 months after 2 nd dose; can only be a Moderna product if 4 years of age or younger; can be either Pfizer-BioNTech or Moderna product if 5 years of age or older <u>Up to Date</u> = immediately after booster
18 years and older	Primary Series	Primary Series; given 4-8 weeks after 1 st dose	Updated Booster; given at least 2 months after 2 nd dose/last booster; can be either Pfizer-BioNTech or Moderna product <u>Up to Date</u> = immediately after booster

Novavax

Age Group	Dose 1	Dose 2	Bivalent Booster
12 years – 17 years	Primary Series	Primary Series; given 3-8 weeks after 1 st dose	Updated Booster; given at least 2 months after 2 nd dose; can be either be Pfizer-BioNTech or Moderna product <u>Up to Date</u> = immediately after booster
18 years and older	Primary Series	Primary Series; given 3-8 weeks after 1 st dose	Updated Booster; given at least 2 months after 2 nd dose; can be either Pfizer-BioNTech or Moderna product <i>*a monovalent Novavax booster is available in limited situations; see CDC website for details</i> <u>Up to Date</u> = immediately after booster

Johnson & Johnson (Janssen)

*this vaccine is authorized for use in certain limited situations; see CDC website for details

Bivalent Booster	Dose 1	Dose 2	Bivalent Booster
18 years and older (not moderately or severely immunocompromised)	Primary Series	N/A	<p>Updated Booster; given at least 2 months after 1st dose/last booster; <i>can be either Pfizer-BioNTech or Moderna product</i></p> <p>*a monovalent Johnson & Johnson/Janssen booster is available in limited situations; see CDC website for details</p> <p><u>Up to Date</u> = immediately after booster</p>
18 years and older (moderately or severely immunocompromised)	Primary Series	Primary Series; given at least 4 weeks after 1 st dose; <i>should be either Pfizer-BioNTech or Moderna product</i>	<p>Updated Booster; given at least 2 months after 1st dose/last booster; <i>can be either Pfizer-BioNTech or Moderna product</i></p> <p>*a monovalent Johnson & Johnson/Janssen booster is available in limited situations; see CDC website for details</p> <p><u>Up to Date</u> = immediately after booster</p>

*If you have completed your primary series, but are not yet eligible for a booster, you are considered Up to Date.

Considerations:

People ages 6 months – 64 years, and especially males ages 12 through 39 years may consider getting the 2nd primary series dose of an mRNA COVID-19 vaccine (Pfizer-BioNTech or Moderna) 8 weeks after the 1st dose. A longer time between the 1st and 2nd primary series doses may increase how much protection the vaccines offer and further minimize the rare risk of myocarditis and pericarditis.

Anyone wanting protection due to high levels of community transmission, people ages 95 years and older, or people who are more likely to get very sick from COVID-19 should get the second dose of Pfizer-BioNTech COVID-19 vaccine 3 weeks (21 days) after the first dose, Moderna COVID-19 vaccine 4 weeks (28 days) after the first dose, or Novavax COVID-19 vaccine 3 weeks (21 days) after the first dose.

For children ages 6 months to 4 years receiving the Pfizer-BioNTech COVID-19 vaccine, the primary series is three doses. For the third dose, the Pfizer-BioNTech bivalent product is given.

CDC does not recommend mixing products for your primary series doses. Children ages 5 years and younger who received a Pfizer-BioNTech primary series must also get a Pfizer-BioNTech booster. Children ages 4 years and younger who received a Moderna primary series must also get a Moderna booster. Children ages 6 years and older who received a Pfizer-BioNTech primary series and children ages 5 years and older who received a Moderna primary series may receive either a Pfizer-BioNTech or Moderna booster regardless of the product used for their primary series. Teens and adults may receive either a Pfizer-BioNTech or Moderna booster regardless of the product used for their primary series. Only the updated (bivalent) mRNA booster is authorized at this time; you can no longer receive the original (monovalent) mRNA booster.

Johnson and Johnson/Janssen COVID-19 vaccine is only authorized for use in certain limited situations. A second primary series dose of Johnson and Johnson/Janssen is recommended for individuals 18 years and older who are moderately or severely immunocompromised.

Johnson and Johnson/Janssen and Novavax monovalent COVID-19 vaccines may be used as a booster in certain limited situations.